
Placental transcriptomes in the common aneuploidies reveal critical regions on the trisomic chromosomes and genome-wide effects.

Journal: Prenat Diagn

Publication Year: 2016

Authors: Katherine Bianco, Matthew Gormley, Jason Farrell, Yan Zhou, Oliver Oliverio, Hannah Tilden, Michael McMaster, Susan J Fisher

PubMed link: 27328057

Funding Grants: SFSU Bridges to Stem Cell Research

Public Summary:

OBJECTIVE: Chromosomal aberrations are frequently associated with birth defects and pregnancy losses. Trisomy 13, Trisomy 18 and Trisomy 21 are the most common, clinically relevant fetal aneusomies. This study used a transcriptomics approach to identify the molecular signatures at the maternal-fetal interface in each aneuploidy. METHODS: We profiled placental gene expression (13-22 weeks) in T13 (n = 4), T18 (n = 4) and T21 (n = 8), and in euploid pregnancies (n = 4). RESULTS: We found differentially expressed transcripts (≥ 2 -fold) in T21 (n = 160), T18 (n = 80) and T13 (n = 125). The majority were upregulated and most of the misexpressed genes were not located on the relevant trisomic chromosome, suggesting genome-wide dysregulation. A smaller number of the differentially expressed transcripts were encoded on the trisomic chromosome, suggesting gene dosage. In T21, <10% of the genes were transcribed from the Down syndrome critical region (21q21-22), which contributes to the clinical phenotype. In T13, 15% of the upregulated genes were on the affected chromosome (13q11-14), and in T18, the percentage increased to 24% (18q11-22 region). CONCLUSION: The trisomic placental (and possibly fetal) phenotypes are driven by the combined effects of genome-wide phenomena and increased gene dosage from the trisomic chromosome. (c) 2016 John Wiley & Sons, Ltd.

Scientific Abstract:

OBJECTIVE: Chromosomal aberrations are frequently associated with birth defects and pregnancy losses. Trisomy 13, Trisomy 18 and Trisomy 21 are the most common, clinically relevant fetal aneusomies. This study used a transcriptomics approach to identify the molecular signatures at the maternal-fetal interface in each aneuploidy. METHODS: We profiled placental gene expression (13-22 weeks) in T13 (n = 4), T18 (n = 4) and T21 (n = 8), and in euploid pregnancies (n = 4). RESULTS: We found differentially expressed transcripts (≥ 2 -fold) in T21 (n = 160), T18 (n = 80) and T13 (n = 125). The majority were upregulated and most of the misexpressed genes were not located on the relevant trisomic chromosome, suggesting genome-wide dysregulation. A smaller number of the differentially expressed transcripts were encoded on the trisomic chromosome, suggesting gene dosage. In T21, <10% of the genes were transcribed from the Down syndrome critical region (21q21-22), which contributes to the clinical phenotype. In T13, 15% of the upregulated genes were on the affected chromosome (13q11-14), and in T18, the percentage increased to 24% (18q11-22 region). CONCLUSION: The trisomic placental (and possibly fetal) phenotypes are driven by the combined effects of genome-wide phenomena and increased gene dosage from the trisomic chromosome. (c) 2016 John Wiley & Sons, Ltd.

Source URL: <http://www.cirm.ca.gov/about-cirm/publications/placental-transcriptomes-common-aneuploidies-reveal-critical-regions>